

Separating strains in metagenomic long-read assemblies

Roland Faure^{1,2}, Jean-François Flot¹, Dominique Lavenier²

¹Université libre de Bruxelles (ULB) - Belgium

²Université de Rennes, IRISA - France

ISMB/ECCB Lyon 2023

Microbiomes are studied at species-level

Identifying Keystone Species in the Human Gut Microbiome from Metagenomic Timeseries Using Sparse Linear Regression

Charles K. Fisher, Pankaj Mehta 

dozens of microbial species could modulate or contribute to cancer
N. Cullin *et al.*, *Microbiome and cancer*, 2021

Metabolic modeling of species interaction in the human microbiome elucidates community-level assembly rules

Bole-Lvay and Eshnar Borenstein  [Authors Info & Affiliations](#)

bacteria are rare. Of the 639 species identified in a population study of 1135 Dutch individuals, 469 (73%) were present in
R.K. Weersma *et al.*, *Interaction between drugs and the gut microbiome*, 2020

difficulties to attribute the species that produce the identified metabolite.
M.S. Afridi *et al.*, *Plant Microbiome Engineering : Hopes or Hypes*, 2022

invasions into soil communities [50]. Similarly, low abundance bacterial species largely contributed to the production of antifungal volatile compounds that protect the plant against soil-borne
S. Compant *et al.* *A review on the plant microbiome : Ecology, functions, and emerging trends in microbial application*, 2019

Microbiomes are studied at species-level

Identifying Keystone Species in the Human Gut Microbiome from Metagenomic Timeseries Using Sparse Linear Regression

Charles K. Fisher, Pankaj Mehta 

dozens of microbial species could modulate or contribute to cancer
N. Cullin *et al.* , *Microbiome and cancer*, 2021

Metabolic modeling of species interaction in the human microbiome elucidates community-level assembly rules

Bole-Ly and Esham Borenstein  [Authors info & Affiliations](#)

bacteria are rare. Of the 639 species identified in a population study of 1135 Dutch individuals, 469 (73%) were present in
R.K. Weersma *et al.* , *Interaction between drugs and the gut microbiome*, 2020


difficulties to attribute the species that produce the identified metabolite.
M.S. Alridi *et al.* , *Plant Microbiome Engineering : Hopes or Hypes*, 2022

invasions into soil communities [50]. Similarly, low abundance bacterial species largely contributed to the production of antifungal volatile compounds that protect the plant against soil-borne
S. Compant *et al.* *A review on the plant microbiome : Ecology, functions, and emerging trends in microbial application*, 2019

Knowledge gaps remain: strain diversity


Ryan Caldwell, Wei Zhou, Julia Oh, *Strains to go: interactions of the skin microbiome beyond its species*, 2022

(Meta)genome assembly: similar regions get collapsed



- ▶ When divergence is small compared to the error rate of reads, it is discarded as sequencing errors

Assembling close strains is difficult



- ▶ Unknown (potentially high) number of strains
- ▶ Uneven coverage
- ▶ One of the reasons microbiomes are studied at species-level

State of the art (long reads)

Article | [Open Access](#) | [Published: 23 July 2021](#)

Strainberry: automated strain separation in low-complexity metagenomes using long reads

[Riccardo Vicedomini](#) ✉, [Christopher Quince](#), [Aaron E. Darling](#) & [Rayan Chikhi](#)

[Nature Communications](#) **12**, Article number: 4485 (2021) | [Cite this article](#)

stRainy: assembly-based metagenomic strain phasing using long reads

Ekaterina Kazantseva, Ataberk Donmez, Mihai Pop, Mikhail Kolmogorov

doi: <https://doi.org/10.1101/2023.01.31.526521>

HairSplitter

- From an assembly, separate the contigs that are present in several versions

```
AACTGTGTCCT-TAGAGCGATTTCGCGACGTA
AACGGTGTCCCTATGGAGCG--TCGCGACCGTA
AACTGTGTCCTATAGAGCGATACGCGACCGTA
AACTGTGTCCT-TAGAGCGATTTCGCGACGTA
AACGGTGTCCCTATAGAGCGATTTCGCGACCGTA
AACGGTGTCCCTATAGAGCGATTTCGCGACCGTA
AACTGTGACCCCTATAGAGCGATACGCGACCGTA
AACTGTGTCCT-TAGAGCGATTTCGACGTA
AACTGCGTCCCTATAGAGCGATACGCGACCGTA
```

Input: All reads, the draft assembly
Output: Strain-separated assembly

```
AACTGTGTCCT-TAGAGCGATTTCGCGACGTA
AACTGTGTCCT-TAGAGCGATTTCGCGACGTA
AACTGTGTCCT-TAGAGCGATTTCGACGTA
```

```
AACGGTGTCCCTATGGAGCG--TCGCGACCGTA
AACGGTGTCCCTATAGAGCGATTTCGCGACCGTA
AACGGTGTCCCTATAGAGCGATTTCGCGACCGTA
```

```
AACTGCGTCCCTATAGAGCGATACGCGACCGTA
AACTGTGACCCCTATAGAGCGATACGCGACCGTA
AACTGTGTCCTATAGAGCGATACGCGACCGTA
```

- Hairsplitter*: One given to hair-splitting or making sophisticated distinctions in reasoning. - *The Century Dictionary*.

Let's try to split the reads

```
Draft assembly  AACTGTGTCCCTATAGAGCGATTTCGCGACCGTACCTCGGAAGCTGAAGTGT
read_1          AACTGTGTCCCT-TAGAGCGATTTCGCGAGCGTATCTCTCGGAAGCTGAA-TGT
read_2          AACGGTGTCCATATGGAGCG--CCGCGACCGTATCTCGAAAGCAGAAAGTGT
read_3          AACTGTGTCCCTATAGAGCGATTTCGCGACCGTACCTCGGAAGCTGAAGTGT
read_4          AACTGTGTCCAT-TAGAGCGATTTCGCGAGCGTATCTCTCGGAAGCTGAA-TGT
read_5          AACGGTGTCCATATAGAGCGATCCGCGACCGTACCTCGAAAGCTGAAGTGT
read_6          AACGGTGTCCCTATAGAGCGATCCGCGACCGTACCTCGAAAGCAGAAATGT
read_7          AACTGTGACCCTATAGAGCGATTTCGCGACCGTACCTCGGAAGCAGAAAGTGT
read_8          AACTGTGTCCAT-TAGAGCGATTTCGCAAGCGTACCTCGGAAGCTGAA-TGT
read_9          AACTGCGTCCCTATAGAGCGATTTCGCGACCGTACCTCGGAAGCAGAAAGTGT
```

First intuition: calling variants


Draft assembly	AACTGTGTCCCTATAGAGCGATTTCGCGACCGTACCTCGGAAGCTGAAGTGT								
read_1	T	C	-		T	G	T	G	T
read_2	G	A	A		C	C	T	A	A
read_3	T	C	A		T	C	C	G	T
read_4	T	A	-		T	G	T	G	T
read_5	G	A	A		C	C	C	A	T
read_6	G	C	A		C	C	C	A	A
read_7	T	C	A		T	C	C	G	A
read_8	T	A	-		T	G	C	G	T
read_9	T	C	A		T	C	C	G	A

First intuition: calling variants

Draft assembly	AACTGTGTCCCTATAGAGCGATTTCGCGACCGTACCTCGGAAGCTGAAGTGT									
read_1	T	C	-		T	G	T	G	T	-
read_2	G	A	A		C	C	T	A	A	G
read_3	T	C	A		T	C	C	G	T	G
read_4	T	A	-		T	G	T	G	T	-
read_5	G	A	A		C	C	C	A	T	G
read_6	G	C	A		C	C	C	A	A	A
read_7	T	C	A		T	C	C	G	A	G
read_8	T	A	-		T	G	C	G	T	-
read_9	T	C	A		T	C	C	G	A	G

- Many of these positions are not actual variants

HairSplitter's key idea: keeping only robust variants




- ▶ A variant is robust if the partition induced is present ≥ 3 times

Separating the reads


1. Compute pairwise distance between reads

	read_1	read_2	read_3	read_4	read_5	read_6	read_7	read_8	read_9
read_1	0.3	0.1	0.9	0.6	0.5	0.0	0.6	0.9	-
read_2	0.6	0.7	0.9	0.0	0.1	0.8	0.6	-	-
read_3	0.1	0.7	0.1	0.9	0.5	0.8	-	-	-
read_4	0.7	0.0	0.9	0.9	0.9	-	-	-	-
read_5	0.8	0.6	0.8	0.1	-	-	-	-	-
read_6	0.9	0.6	0.8	-	-	-	-	-	-
read_7	0.0	0.4	-	-	-	-	-	-	-
read_8	0.7	-	-	-	-	-	-	-	-
read_9	-	-	-	-	-	-	-	-	-

2. Build graph using KNN




3. Cluster graph



Re-assemble reads

Reads aligned on
draft assembly

```
AACTGTGTCCT-TAGAGCGATTTCGCGAGCGTA
AACGGTGTCCCTATGGAGCG--TCGCGACCGTA
AACTGTGTCCTATAGAGCGGATACGCGACCGTA
AACTGTGTCCT-TAGAGCGATTTCGCGAGCGTA
AACGGTGTCCCTATAGAGCGATTTCGCGACCGTA
AACGGTGTCCCTATAGAGCGATTTCGCGACCGTA
AACTGTGACCCCTATAGAGCGATACGCGACCGTA
AACTGTGTCCT-TAGAGCGATTTCGCAAGCGTA
AACTGCGTCCCTATAGAGCGATACGCGACCGTA
```




Simulated data


- ▶ Mix of 2 to 10 *E. coli* RefSeq genomes. Simulated reads
- ▶ Metrics: metaQuast completeness, error rate, **31-mer completeness...**

Simulated data

- ▶ Mix of 2 to 10 *E. coli* RefSeq genomes. Simulated reads
- ▶ Metrics: metaQuast completeness, error rate, **31-mer completeness**...



(a) Nanopore sequencing (12% error), 50x per strain



(b) HiFi sequencing, 10x per strain

- ▶ HairSplitter handles very well high number of strains

Mock data

- Zymobiomics gut microbiome standard: contains a mix of 5 *E. coli* strains

	metaFlye	metaFlye+Strainberry	metaFlye+HairSplitter
Nanopore Q9	0.586	0.749	0.957
Nanopore Q20	0.7524	0.9527	0.961
PacBio HiFi	0.9589	0.9793	0.9895

Table: 31-mer completeness of assemblies w.r.t. the reference


Real data

- ▶ 5 *Vagococcus fluvialis* strains sequenced with Nanopore **barcoded** reads (doi.org/10.1186/s12864-022-08842-9).

	metaFlye	metaFlye+Strainberry	metaFlye+HairSplitter
Nanopore	0.718	0.7398	0.9042

Table: 31-mer completeness of assemblies w.r.t. a Flye assembly where reads from different strains were separated.

Limitation: Contiguity



- Strains are separated only locally
- N50 can decrease significantly

Take-home message

- ▶ HairSplitter **reconstruct collapsed sequences** from “draft” assemblies obtained by any means
- ▶ HairSplitter also works well on **complex metagenomes** and with **high-error rate** reads

Take-home message


- ▶ HairSplitter **reconstruct collapsed sequences** from “draft” assemblies obtained by any means
- ▶ HairSplitter also works well on **complex metagenomes** and with **high-error rate** reads
- ▶ Available **prototype** ! github.com/RolandFaure/HairSplitter

Acknowledgements


- ▶ Dominique Lavenier and Jean-François Flot for their supervision
- ▶ Rumen Andonov and Tam Truong for their help in formalizing the problem
- ▶ The EEB-EBE and GenScale teams




Behaviour of HairSplitter/BiHap: coverage



Local strain separation



Local strain separation



metaQuast evaluation

Genome statistics	flye	strainberry	hairstripper
+ Genome fraction (%)	89.353	58.457	97.891
+ Duplication ratio	1.228	1.429	2.656
+ Largest alignment	330 446	312 915	272 305
+ Total aligned length	4 053 565	5 347 608	10 043 014
+ NGA50	...	...	...
+ LGA50	...	...	...
Misassemblies			
+ # misassemblies	50	51	141
+ Misassembled contigs length	2 412 308	3 237 621	5 265 749
Mismatches			
+ # mismatches per 100 kbp	1914.71	766.25	1887.03
+ # indels per 100 kbp	1488.89	1170.19	1274.4
+ # N's per 100 kbp	0	135.18	0
Statistics without reference			
+ # contigs	240	295	399
+ Largest contig	845 747	551 381	611 375
+ Total length	4 573 883	5 693 921	10 384 735
+ Total length (>= 1000 bp)	4 552 674	5 669 228	10 369 591
+ Total length (>= 10000 bp)	3 977 976	4 941 261	9 534 633
+ Total length (>= 50000 bp)	3 016 695	3 786 882	7 396 612

[Extended report](#)

metaQuast evaluation of
Vagococcus fluvialis assemblies

Genome statistics	flye	strainberry	hairstripper
+ Genome fraction (%)	80.324	95.595	95.321
+ Duplication ratio	1.133	1.976	2.002
+ Largest alignment	233 623	236 712	254 460
+ Total aligned length	7 186 249	16 406 302	17 127 330
+ NGA50	...	...	...
+ E_coli_ref	-	26 028	12 000
+ LGA50	...	...	...
Misassemblies			
+ # misassemblies	66	74	99
+ Misassembled contigs length	4 083 597	4 961 648	4 365 176
Mismatches			
+ # mismatches per 100 kbp	1771.49	1684.86	1474.75
+ # indels per 100 kbp	163.72	114.67	88.75
+ # N's per 100 kbp	0	40.1	0
Statistics without reference			
+ # contigs	1347	1814	7265
+ Largest contig	4 733 384	4 799 087	2 026 315
+ Total length	60 410 911	69 375 520	71 918 465
+ Total length (>= 1000 bp)	60 402 812	69 334 461	71 856 894
+ Total length (>= 10000 bp)	58 032 898	66 014 854	52 120 366
+ Total length (>= 50000 bp)	39 735 723	44 207 867	31 223 699

[Extended report](#)

metaQuast evaluation of Zymobiomics
gut microbiome standard Nanopore Q20
assemblies